

Pathology of the central nervous system.

Pathology of the central nervous system.

I. Microspecimens:

№ 68. Ischemic infarction in the brain. (*gray softening*). (H.E. stain).

Indications:

1. Focus of necrosis with cerebral tissue rarifying.
2. Adjacent cerebral tissue with edema.

In microspecimen there is an area of thinning of brain tissue, most nerve cells are anucleated, the remaining cells are with signs of karyopyknosis. Pronounced perivascular and pericellular edema, perifocal leukocytic infiltration, in some places proliferation of glial elements (reactive gliosis) and leukocytic infiltration can be seen. Arterioles are with thickened, sclerosed walls, with signs of hyalinosis (hyaline arteriolosclerosis).

Cerebral ischemic infarction is associated in the vast majority of cases with thrombosis or thromboembolism of the cerebral arteries, which causes hypoperfusion or even interruption of arterial blood supply in a certain area of the brain. It is one of the manifestations of cerebrovascular disease. Cerebral artery thrombosis is found in atherosclerosis, in states of hypercoagulability. Thromboembolism can occur in atherosclerosis of the carotid arteries and aorta, in atrial fibrillation. There are cases of cancerous embolism (eg., in cardiac myxoma), vegetation in the left cavities of the heart in infectious endocarditis, in myocardial infarction, cardiomyopathies. Macroscopically, the infarct area has an irregular shape, imprecise boundaries, the border between gray and white substance is faded, flabby consistency, whitish-gray color (white or gray softening). Clinically it is manifested by paralysis, aphasia. Consequences: macrophage resorption of necrotic masses and fibroglial organization, cystic transformation.



№ 68. Ischemic infarction in the brain. (gray softening). (H.E. stain).


№ 28. Pyogenic leptomeningitis. (*H-E. stain*).

Indications:

1. Neutrophilic infiltration of leptomeninges.
2. Edematous cerebellar cortex.

In microspecimen, with naked eye, thickened leptomeniges of intense basophilic color, and cerebellar cortex of eosinophilic color are observed. At the small magnification, leptomeninges is diffuse infiltrated by neutrophilic leukocytes with presence of edema, dilated blood vessels, hyperemia and leukocytic agglomerations in subarachnoid space. In cerebellar cortex there are perivascular and pericellular edema (colorless spaces around vessels and cells), dilation and hyperemia of blood vessels, punctiform hemorrhages and focal infiltrates with neutrophilic leukocytes.

Pyogenic leptomeningitis is an example of phlegmonous purulent inflammation - inflammation without precise delimitation, in which the exudate is diffusely spread between the tissue elements. The pus spreads along the intermuscular spaces, adipose tissue, neuro-vascular trunks, etc. In addition to leptomeninges, it is encountered in adipose tissue, muscles, walls of cavitory and tubular organs (vermicular appendix, gallbladder, stomach, intestine). The most common causative factor of pyogenic leptomeningitis is meningococcus. As a result, resorption of the exudate and complete resolution may develop, or thickening of the meningeal membranes may occur and adhesions between the membranes and between them and brain surface, which favors the appearance of cystic cavities in the thickness of the leptomeninges or even of the internal hydrocephalus caused by the stenosis or obstruction of the Magendie and Luschka holes.



№ 28. Pyogenic leptomeningitis. (H-E. stain).


№ 45. Glioblastoma multiforme. (H.E. stain).

Indications:

1. Atypical and polymorphic tumoral cells.
2. Focus of necrosis.

Tumor tissue has a rich, dense cellularity, marked cellular and nuclear polymorphism, spindle-shaped cells, small polymorphic cells, giant cells, some polynucleate, mitotic figures, eosinophilic anucleate foci of coagulative necrosis, around which the tumor cells are arranged in palisade. Numerous vessels are observed, some with intraluminal thrombosis, others stenosed by endothelial hyperplasia. Foci of hemorrhages and necrosis are seen as well.

CNS tumors account for 2% of the mortality rate of malignant tumors in adults and 20% - in children. Glioblastoma is found in 40% of all primary brain tumors in adults. It is an aggressive tumor, of glial origin, with low differentiation. It may occur de novo or develop through the progression of diffuse astrocytoma. The local effects consist of compression and destruction of adjacent brain tissue, infarctions and hemorrhages by invasion of blood vessels, peritumoral edema, disorders of cerebrospinal fluid circulation. It metastasizes within the CNS and does not produce extracranial metastases (or patients die before such metastases occur).



№ 45. Glioblastoma multiforme. (H.E. stain).


№ 123. Transitional meningioma (H.E. stain).

Indications:

1. Nests of cells with a spiral appearance.
2. Psammoma bodies.

The tumor consists of concentric of concentric agglomerations of elongated syncytial and fibroblastic cells, among which multiple psammoma bodies are determined.

Meningioma is a benign tumor, usually solitary, which occurs predominantly in women in the sixth decade of life, accounting for about 20% of all intracranial tumors. It is derived from the meningotheial cells of the arachnoid membrane (arachnoid cells). It is most commonly located on the parasagittal line along the falx cerebri. Macroscopically, it represents tumor node with a diameter of 1-10 cm, attached to the dura mater, which compresses the brain tissue, but does not invade it, has a dense consistency, on section fibrous aspect, there may be foci of intratumoral calcinosis. The most common histological variants are: meningotheial, fibrous (fibroblastic) and transitional (with mixed pattern of meningotheial and fibroblastic meningioma). Clinical symptoms depend on the location and size of the tumor. Very rarely it can become malignant.



№ 123. Transitional meningioma (*H.E. stain*).


II. Macrospecimens:

№ 121. Brain hematoma.

In the brain there is an accumulation of dark red coagulated blood (hematoma), the adjacent brain tissue is softened, flabby, edematous.

The main cause of parenchymal cerebral hemorrhage is rupture of the arteries. It is one of the manifestations of cerebrovascular disease. It is most commonly observed in hypertension (formation of microaneurysms, fibrinoid necrosis of arterioles in the hypertensive crisis), leukemias, severe thrombocytopenia, congenital aneurysms. The most common location is in the basal ganglia and thalamus - 65%, pons Varolii - 15%, cerebellum - 10%. Hemorrhage causes both direct tissue damage and secondary ischemic damage. Clinically it is manifested by paralysis, aphasia.

Consequences: fibroglial organization, cystic cavities.



№ 121. Brain hematoma.


№ 122. Brain tumor - glioblastoma.

On the section of the brain there is a tumor node with a diameter of several cm, irregular shape, without clear boundaries, white-gray color, with foci of necrosis, hemorrhages, small cystic cavities, located in the white matter, adjacent brain tissue is with signs of rarifying and softening. (microspecimen nr. 45).

№ 123. Hydrocephalus.

The brain is enlarged, the lateral ventricles considerably dilated, the brain tissue is atrophied.

Hydrocephalus - excessive accumulation of cerebrospinal fluid in the ventricular system, the cause being stenosis of the foramina of Monro, Sylvius aqueduct, the Magendie and Luschka holes. It can develop as a result of leptomeningitis, brain tumors, CNS trauma, hydrocephalus can be ex vacuo, when compensatory hyperproduction of cerebrospinal fluid occurs as a reaction to decreased brain tissue in Alzheimer's disease, after parenchymal hemorrhage, cerebral infarction. If the entire ventricular system is dilated, hydrocephalus is called communicating, and if there is an obstacle inside the ventricular system -it is called non-communicating hydrocephalus, in such cases only a portion of the ventricle is dilates. The non-communicating variant can be found in ependymal tumors, in the choroid plexus papilloma, cerebral abscesses, parenchymal, intraventricular, epi- or subdural hemorrhages. Hydrocephalus causes compression atrophy of the brain parenchyma.



№ 122. Brain tumor - glioblastoma.



№ 123. Hydrocephalus.


№ 124. Purulent leptominengitis.

The leptomeninges on the convex surface of the brain, especially in the frontal, temporal and parietal areas, is thickened, edematous, imbued with purulent exudate of gray-yellow color, the inflamed area has no clear boundaries blood vessels are dilated and hyperemic.





Purulent leptomeningitis is of bacterial origin, in adolescents and young adults it is caused more frequently by meningococcus (Neisseria meningitidis), and in the elderly - by streptococci. In some cases meningitis develops secondarily by spreading the infection from adjacent purulent foci (in mastoiditis, otitis media, paranasal sinusitis) or by hematogenous dissemination in infectious endocarditis, pneumonia. Morphologically it is manifested by diffuse purulent inflammation (phlegmonous) of the meningeal membrane, the pus also appears in the subarachnoid space, in the grooves, the cerebrospinal fluid becomes cloudy, contains abundant neutrophilic leukocytes, can be associated meningoencephalitis, ventriculitis, brain abscesses. Consequently, exudate resorption and complete resolution may occur or residual morphological changes may occur, e.g., thickening of the meningeal membranes and the formation of adhesions both between the membranes and between them and the surface of the brain, which favors the appearance of cystic cavities in the subarachnoid space or even internal hydrocephalus.



№ 124. Purulent leptomeningitis.



**Brain atrophy
Alzheimer's disease.**



Alzheimer's disease:
a,b - senile plaques,
c – neurofibrillary tangle (*silver impregnation*),
d – amyloid angiopathy (*Congo red stain*).



Parkinson's disease. Damage to the black substance.





Parkinson's Disease

⇒ Signs



- ⇒ Akinesia (RT, initiation)
- ⇒ Bradykinesia (MT, slow)
- ⇒ Rigidity
- ⇒ Tremor
- ⇒ Postural instability, difficulty multi-tasking, sleep difficulties, cognitive deficits.

⇒ Cause


- ⇒ Denervation of DAergic neurons in SNpc
- ⇒ Loss of DA in striatum




Boala Parkinson
Manifestări clinice.



Huntington's Disease Affects the Brain's Basal Ganglia



Autosomal Dominant Inheritance




Normal


ALS



Amyotrophic lateral sclerosis,
atrophy of the anterior motor nerve
roots of spinal cord.



Multiple sclerosis, plaques
(foci of demyelination).







**Purulent bacterial
leptomeningitis
and brain abscess.**



Cerebral edema with herniation of the cerebellar tonsils in the occipital foramen magnum.

**Intracerebral
hematomas.**







a

Cerebral infarction :




*a – hemorrhagic infarction,
b, c – ischemic infarction*







b





c





**a, b – ischemic softening with cystic transformation;
c,d – lacunar infarction.**



Glioblastoma multiforme.





Meningioma.





**Ependymoma in the area
of IV ventricle.**





**Carcinoma metastasis
in the brain.**

CNS

- **Normal**

- Neurons

- Glia

- Astrocytes

- Oligodendrocytes

- Ependymal Cells

- Microglia

- **Pathology (13 Questions)**


Classical Disease Patterns

- **Degenerative**
- **Inflammatory**
- **Neoplastic**

Classical **CNS** Disease Patterns

- **Degenerative**
- **Inflammatory**
- **Neoplastic**
- **Traumatic**

- 1) What are general patterns of CNS **cell** pathology?
 - 2) What are the consequences of ↓ ↑ CNS **pressure**?
 - 3) What are common patterns of CNS **malformations**?
 - 4) What are common **perinatal** CNS injuries?
 - 5) What are the patterns of CNS **trauma**?
 - 6) What are the patterns of CNS **vascular** disease?
 - 7) What are the patterns of CNS **infection**?
-
- 8) What are the patterns of CNS **prion** disease?
 - 9) What are the patterns of CNS **demyelinating** disease?
 - 10) What are the patterns of CNS **degenerative** disease?
 - 11) What are the CNS **genetic metabolic** diseases?
 - 12) What are the CNS acquired **metabolic/toxic** diseases?
 - 13) What are the CNS **tumors**?



Ribosomes

Rough ER

Membrane

Microtubules

Mitochondrion

Nucleus


Axon hillock



Axon



Smooth ER





Golgi apparatus


Polyribosome























CELLULAR REACTIONS


•Neurons

- Acute (**RED** neuron, karyolysis)
- Subacute, chronic, cell loss, gliosis
- Axonal
- Inclusions (lipid, prot., carb., viruses)

•Glia, “gliosis”


- Swelling
- Fibers

- Inclusions



ACUTE NEURONAL INJURY

“**RED**” NEURONS



CEREBRAL EDEMA


(normal weight 1200-1300 grams)

- **Vasogenic** (disrupted BBB)
 - Intravascular → INTER-cellular
- **Cytotoxic**
 - INTRA-cellular

CEREBRAL EDEMA

- Subfalcine (**SUPRA-tentorial**)
- Cingulate (**TENTORIAL**)
- Cerebellar tonsillar (**SUB-tentorial, or INFRA-tentorial**)






Subfalcine
herniation

Transtentorial
herniation

Tonsillar herniation







CEREBRAL EDEMA


- **DDX:**

- **EVERYTHING**

- **SYMPTOMS**

- **HEADACHE**
- **HALLUCINATIONS**
- **COMA**
- **DEATH**


HYDROCEPHALUS




HYDROCEPHALUS

- **Impaired RESORPTION**
- Increased PRODUCTION
- **OBSTRUCTION**
- **COMMUNICATING (entire)**
- **NON-COMMUNICATING (part)**
- **HIGH Pressure**
- **NORMAL Pressure**







Aqueductal stenosis








CM



CNS MALFORMATIONS

- **Neural Tube**
 - **Anencephaly, Encephalocele, Spina Bifida**
- **Forebrain**
 - **Polymicrogyria, Holoprosencephaly, Agenesis of Corpus Callosum**
- **Posterior Fossa (Infratentorial)**
 - **Arnold Chiari (infratentorial herniation), Dandy-Walker (cerebellar cyst)**
- **Syringomyelia/Hydromyelia**



SPINA BIFIDA








POLYMICROGYRIA



HOLOPROSENCEPHALY




Corpus callosum



SYRINGOMYELIA




(note **“SYRINX”**)



PERINATAL Brain Injuries

- Intraparenchymal Hemorrhage
- Intraventricular hemorrhage (premies)
- Periventricular “leukomalacia” (i.e., infarcts)

- Cerebral “Palsy” refers to nonprogressive diffuse cerebral pathology apparent at childbirth




CNS TRAUMA

- Skull Fractures
- Parenchymal Injuries
- Traumatic Vascular Injury
- Sequelae
- Spinal Cord Trauma


BRAIN TRAUMA

- Contusion (bruise)
- Laceration (tear)
- Coup/Contre-Coup
- Concussion





“HAIRLINE”




**“DEPRESSED”,
aka**

“DISPLACED”


HEMATOMAS/HEMORRHAGE



- **EPIDURAL (fx)**
- **SUBDURAL (trauma NO fx)**
- **SUBARACHNOID (arterial, no trauma)**
- **INTRAPARENCHYMAL (any)**
- **INTRAVENTRICULAR (no trauma, rare in adults, common in premies)**






EPIDURAL HEMATOMA







SUBDURAL HEMATOMA




SUBARACHNOID



INTRAPARENCHYMAL



INTRAPARENCHYMAL



INTRAVENTRICULAR

CNS TRAUMA SEQUELAE


- **Hydrocephalus (WHY?)**
- **Dementia (Punch Drunk Syndrome)**
- **Diffuse Axonal Injury (white matter)**

89 ME 128



SPINAL CORD TRAUMA

- Parallels BRAIN patterns of injury on a cellular basis
- Usually secondary to spinal column displacement
- Level of injury mirrors motor loss:
Death → Quadriplegia → Paraplegia




Cerebrovascular Diseases (CVA, “Stroke”)

• Ischemic (↓ blood and O₂)

- Global
- Focal (regional):
 - **ACUTE**: edema → neuronal microvacuolization → pyknosis → karyorrhexis → neutrophils
 - **CHRONIC**: macrophages → gliosis

• Hemorrhagic (rupture of artery/aneurysm)

Anterior Cerebral A.



Middle Cerebral A.

Posterior Cerebral A.




THROMBOTIC MCA







HEMORRHAGIC



ACA





A) EDEMA

B) "RED" NEURONS

C) POLYs

D) MONO's (MACs)

E) GLIOSIS


**Histopathologic
progression of
CNS infarcts**

HYPERTENSIVE CVA

- **Intracerebral**


- **Basal Ganglia Region**

(lenticulostriate arteries of internal capsule, putamen)






HYPERTENSIVE CVA



LACUNAR INFARCTS




52/98

A coronal section of a human brain, showing the cerebral cortex with its characteristic gyri and sulci. The brain is bisected vertically, revealing the internal structures. In the central region, specifically in the basal ganglia area, there are several small, dark, slit-like hemorrhages. These hemorrhages are located symmetrically on both sides of the midline, near the lateral ventricles. The surrounding brain tissue appears normal in color and texture.


“SLIT” HEMORRHAGE(s)

SUBARACHNOID HEMORRHAGE

- Rupture of large intracerebral arteries which are the primary branches of the anatomical circle (of Willis)
- **Congenital** (“berry” aneurysms)
- **Atherosclerotic** (atherosclerotic aneurysms, or direct wall rupture)







HYPERTENSIVE ENCEPHALOPATHY

• ACUTE

- Headaches
- Confusion
- Anxiety
- Convulsions

• CHRONIC

- Dementia (MID, **M**ulti-**I**nfarct-**D**ementia)
- Gait Disturbances
- Basal Ganglia symptoms

CNS INFECTIONS

- ACUTE MENINGITIS
- ACUTE FOCAL SUPPURATIVE
- CHRONIC BACTERIAL
- VIRAL
- FUNGAL
- OTHER

INFECTIONS

- **Meningitis (generally* bacterial)**

- E. coli, Strep B (neonates)
- H. influenzae (children)
- Neisseria meningitidis (adults)
- Strep. pneumoniae, Listeria (elderly)
- PMNs in CSF, INCREASED protein, REDUCED glucose

- **Encephalitis (generally viral)**


- Arboviruses, HSV, CMV, V/Z, polio, rabies, HIV
- Lymphs and macrophages in perivascular “Virchow-Robbins” spaces


- **Meningoencephalitis**

*

viral, chemical, tumoral








ACUTE FOCAL SUPPURATIVE CNS INFECTIONS


• CEREBRAL ABSCESSSES

- Local (mastoiditis, sinusitis)
- Hematogenous (tooth extraction, sepsis)
- Staph, Strep
- Often fibrous capsule, liquid center


• SUBDURAL EMPYEMA (IN SINUSITIS)

• EXTRADURAL ABSCESS (IN OSTEOMYELITIS)





A45-603






SUBDURAL EMPYEMA

Fat in the epidural space

Epidural venous plexus

Dura mater



Posterior spinal arteries




Abscess in epidural space

CHRONIC BACTERIAL Meningo-encephalitis

- **TB**, brain and meninges
- **SYPHILIS**, gummas in brain
- **LYME DISEASE** (Neuro-Borreliosis)





TUBERCULOMA





VIRAL

Meningo-encephalitis


- ARBO VIRUSES (West Nile, Equines, Venez., many more)
- HSV1
- HSV2
- V/Z
- CMV
- POLIO
- RABIES
- HIV
- Progressive Multifocal Leukoencephalopathy (JC)
- Subacute Sclerosing Panencephalitis (Measles)






**VIRAL
ENCEPHALITIS
PERIVASCULAR
LYMPHOCYTIC
“CUFFING”**





Bitemporal encephalitis is HSV until proven otherwise!



HSV = TEMPORAL lobe(s)






**PERIVASCULAR
GIANT CELLS
in WHITE MATTER in
HIV
ENCEPHALITIS**

PROGRESSIVE MULTIFOCAL LEUKOENCEPHALOPATHY (PML)


- JC Polyoma virus is the cause
- Primarily affects oligodendrocytes
- Ergo, demyelination is the main feature




46

NETX







A





B

PML

SUBACUTE SCLEROSING PANENCEPHALITIS (SSPE)


- VERY rare since measles eradicated
- Thought to be caused by measles virus





FUNGAL MENINGO-ENCEPHALITIS

- **CRYPTOCOCCUS**
- **CANDIDA**
- **ASPERGILLIS**
- **MUCOR**

(Mostly in immunocompromised hosts)






**CRYPTOCOCCUS
MICROABSCESSSES**


OTHERS

- **MALARIA**
- **TOXOPLASMOSIS (in HIV)**
- **AMEBIASIS**
- **TRYPANOSOMES**
- **RICKETTSIAE**
- **ECHINOCOCCUS**



CNS

II



- 1) What are general patterns of CNS **cell** pathology?
- 2) What are the consequences of ↓↑ CSF **pressure**?
- 3) What are common patterns of CNS **malformations**?
- 4) What are common **perinatal** CNS injuries?
- 5) What are the patterns of CNS **trauma**?
- 6) What are the patterns of CNS **vascular** diseases?
- 7) What are the patterns of CNS **infection**?

- 8) What are the patterns of CNS **prion** diseases?
- 9) What are the patterns of CNS **demyelinating** diseases?
- 10) What are the patterns of CNS **degenerative** diseases?
- 11) What are the CNS **genetic metabolic** diseases?
- 12) What are the CNS **acquired metabolic/toxic** diseases?
- 13) What are the CNS **tumors**?

PRION DISEASES

• Creutzfeldt-Jakob Disease (CJD)


- Gerstmann-Straussler-Scheinker syn. (GSS)
- Fatal familial insomnia
- Kuru, human variety (cannibalism)
- Scrapie (sheep and goats)
- Mink transmissible encephalopathy
- Chronic wasting disease (deer and elk)
- **Bovine Spongiform Encephalopathy (BSE)**

PRION DISEASES:

common features

- Infectious agents with apparently no DNA
- DEMENTIA
- Prion Protein (**PrP**) accumulation
- **“SPONGIFORM”** changes in neurons and glia
- TRANSMISSIBLE, FATAL, NO Rx

PRION PROTEIN




Normally found in humans

Exact structure known, 208 amino acids

Specific chromosome, #20, specific genes also known

Requires a **conformational change** to accumulate and do damage



CJD (Creutzfeldt-Jakob)

- 1 per million incidence, 7th decade
- Sporadic cases, **not** epidemic
- Transmitted!
- Familial cases well documented
- Rapidly progressive dementia
- Grey Matter
- Cerebellar ataxia also, usually
- FATAL, no treatment known, like ALL prion diseases


DEMYELINATING DISEASES

- MS (**M**ULTIPLE **S**CLEROSIS)


- MS variants
- ACUTE **D**ISSEMINATED **E**NCEPHALOM**M**YELITIS (ADEM)
- ACUTE **N**ECROTIZING **H**EMORRHAGIC **E**NCEPHALOM**M**YELITIS (ANHE)
- Many, many, many others. Remember: DEMYELINATION is a **NON-SPECIFIC** reaction to MANY types of CNS injury, and demyelination also causes edema

MS


- Cause: ?
- USA prevalence: 1:1000
- F>>M, Ages: 30's, 40's
- Immune response primarily against CNS myelin (white matter)
- Regional area of white matter demyelination is called **"PLAQUE"**
- Increased CSF gamma globulin, i.e., **oligoclonal bands**
- **Often presents with VISUAL problems**
- **EXACERBATIONS/REMISSIONS**




A




B




C






D





PLAQUES, MS



CNS DEGENERATIVE DISEASES

- CORTEX (**dementias**)
- BASAL GANGLIA and BRAIN STEM (**parkinsonian**)
- SPINOCEREBELLAR (**ataxias**)
- MOTOR NEURONS (**muscle atrophy**)

CNS DEGENERATIVE DISEASES




- **CORTEX (dementias)**
 - **ALZHEIMER DISEASE**
 - Frontotemporal
 - **Pick Disease** (also primarily frontal)
 - **Progressive Supranuclear Palsy (PSP)**
 - **CorticoBasal Degeneration (CBD)**
 - **Vascular Dementias (MID)**



ALZHEIMER DISEASE

- Commonest cause of dementias (majority)
- Sporadic, 5-10% familial
- **CORTICAL (grey matter) ATROPHY**
- **NEURITIC PLAQUES*** (extraneuronal)
- **NEUROFIBRILLARY TANGLES**
(intraneuronal)
- **AMYLOID!!!** (i.e., “BETA” amyloid)













Neuritic plaques


Neuritic plaques, stained with
anti- **beta amyloid**
immunostain











Tangle

Nucleus

1 μ m

OTHER CORTICAL DEMENTIAS

(tau gene/protein, tau-opathies)

- FRONTOTEMPORAL
- PICK DISEASE (LOBAR ATROPHY)
- PROGRESSIVE SUPRANUCLEAR PALSY (PSP)
- CORTICOBASAL DEGENERATION (CBD)

- VASCULAR DEMENTIA (MID)

VASCULAR DEMENTIA

- Associated with multiple infarcts, hence the name MID (**M**ultiple **I**nfarct **D**ementia)
 - Lacunar infarcts
 - Cortical microinfarcts
 - Multiple embolic infarcts
- **SECOND** commonest form of dementia after Alzheimer

SC2
IR-TSE/M
SL13

A

10 cm



FH 36 head

CNS DEGENERATIVE DISEASES

- **BASAL GANGLIA and BRAIN STEM**

- **Parkinsonism**

- Parkinson Disease
- Multiple System Atrophy
- Huntington Disease

Parkinsonism

- Is a clinical “syndrome”, NOT a disease
 - Diminished facial expression
 - Stooped posture
 - Slowness of voluntary movement
 - “Festinating” gait (short, fast)
 - Rigidity (cogwheel)
 - “Pillrolling” tremor
- The above clinical findings involve pathology of the SUBSTANTIA NIGRA, and include:

- **PARKINSON DISEASE**


- MULTIPLE SYSTEM ATROPHY
- POSTENCEPHALIC PARKINSONISM
- Progr. Supranuc. Palsy, Cort. Basal Degen. (cortical disorders)

PARKINSON DISEASE

- **PALLOR** of the
SUBSTANTIA NIGRA (and
LOCUS COERULEUS)
- **LEWY BODIES** (alpha-synuclein protein)










LOCUS COERULEUS* in PONS (CERULEUS**)

* 254,000

** 76,000









PARKINSON DISEASE

- **Parkinsonism symptoms, i.e.,**
 - cogwheel rigidity
 - intention tremor
- **Progressive**
- **Hallucinations**
- **Dementia**
- **Symptomatic response to L-DOPA**

MULTIPLE SYSTEM ATROPHY

- MSA
- WIDE SPECTRUM of diseases
- GLIAL CYTOPLASMIC INCLUSIONS (GCIs)
in oligodendrocytes (alpha synuclein)
- Clinically,
 - parkinsonism symptoms
 - autonomic dysfunction



HUNTINGTON DISEASE

- **Classical familial, genetic disease**
- **Progressive motor loss and dementia**
- **“chorea”, i.e. “jerky” movements**
- **Progressive, fatal**
- **Atrophy of basal ganglia, i.e., corpus striatum**



Cortical (basal ganglia) atrophy

Ventricular enlargement

CNS DEGENERATIVE DISEASES

•SPINOCEREBELLAR DEGENERATIONS (ATAXIAS)

- Spinocerebellar ataxias
- Friedrich Ataxia
- Ataxia-Telangiectasia

SPINOCEREBELLAR DEGENERATIONS

- Cerebellar cortex
- Spinal cord
- Peripheral nerves

- FEATURES:

- **ATAXIA** (loss of extremity muscle coordination)

- **SPASTICITY**

- **NEUROPATHIES**

CNS DEGENERATIVE DISEASES

- **MOTOR NEURONS**

- **ALS** (Amyotrophic Lateral Sclerosis, i.e., Lou Gehrig's disease)

- BulboSpinal Atrophy (Kennedy Syndrome)

- Spinal Muscular Atrophy

Amyotrophic Lateral Sclerosis

- Unknown etiology
- Progressive muscle atrophy due to motor neuron loss (lower, upper)
- 5-10% familial
- **Lou Gehrig** had it, so does **Steven Hawking**
- Hand weakness → diaphragm
- Anterior horn cells reduced and gliotic



THE RECORD-BREAKING BESTSELLER NOW IN PAPERBACK

A BRIEF HISTORY OF TIME



From the Big Bang to Black Holes

'This book marries a child's wonder to a genius's intellect. We journey into Hawking's universe, while marvelling at his mind'
Sunday Times




Introduction by Carl Sagan

STEPHEN HAWKING



A.L.S., DEMYELINATION IN CORTICOSPINAL TRACTS



ALS, pathologic changes in anterior horn cells

GENETIC METABOLIC DISEASES

- **NEURONAL STORAGE DISEASES**

- (classical autosomal recessive enzyme deficiencies)

- **“LEUKO”-DYSTROPHIES**

- (abnormal “myelin” synthesis)

- **MITOCHONDRIAL ENCEPHALOPATHIES**


- (mitochondrial gene mutations)


LEUKODYSTROPHIES

- Krabbe
- Metachromatic-
- Adreno-
- Pelizaeus-Merzbacher
- Canavan

ACQUIRED TOXIC/METABOLIC CNS DISEASES

- Vitamin **B1** deficiency (Wernicke-Korsakoff)
- Vitamin **B12** deficiency (vibratory sense)
- **Diabetes** Increased/Decreased GLUCOSE
- Hepatic Failure (**NH₄⁺**)
- **CO** (Cortex, hippocampus, Purkinje cells)
- **CH₃-OH**, Methanol (Retinal ganglion cells)
- **CH₃-CH₂-OH** (acute/chronic, direct/nutrit'l)
- **Radiation** (Brain MOST resistant to Rad. Rx.)
- **Chemo** (Methotrexate + Radiation)





128 Hz

CNS TUMORS

- GLIOMAS (do not metastasize out of the CNS)
 - Astrocytes (I, II, **III, IV**)
 - Oligodendroglioma
 - Ependymoma
- NEURONAL (neuroblastoma)
- POORLY DIFFERENTIATED (medulloblastoma)
- MENINGIOMAS
- LYMPHOMAS
- METASTATIC

CNS TUMORS

• SYMPTOMS?

- Headache
- Vomiting
- Mental Changes
- Motor Problems
- Seizures
- Increased Intracranial Pressure
- **ANY** localizing CNS abnormality

CNS TUMORS


- History
- Physical
- Neurologic exam
- LP (including cytology)
- CT
- MRI
- Brain angiography
- Biopsy


CNS TUMORS


- Benign? Malignant?, Primary vs. met?
- Location?
- Age?
- X-ray Density? MRI signals?
- Calcifications?
- Vascularity?
- Necrosis?
- Liquefaction?
- Edema?
- Compression of neighbors?


GLIOSIS vs. GLIOMA

- Age?
- White vs. Grey Matter?
- Gross texture?
- Vascularity?
- Mitoses?
- (N/C, Pleomorphism, Hyperchromasia)
- Calcifications?
- Cysts?
- Satellitosis?
- Delineation?









68Y
1:05
01-JUN-1998
IMAGE 126
STUDY 7


H-SP VB31
+ : F A






el 90
R
SAT
R 600.0
E 12.0/1
A 03:21


SP -0.5
SL 4.0
FoV 158*210
165 *2560
Cor


**NON
ASTROCYTIC
GLIOMAS**
























Zoom
Me!







MENINGIOMAS

- Occur where dura is
- Very vascular
- BENIGN, but.....(can be damned invasive)
- Can invade skull, etc.
- Only invade (displace) brain in areas adjacent to dura, i.e., parasagittal, falx, tentorium, venous sinuses
- Small, firm, and well defined like a SUPERBALL
- Often (usually?) have **PSAMMOMA** bodies









HIV

METASTATIC CNS TUMORS

- **LUNG**

- BREAST

- MELANOMA

- KIDNEY

- GI

“PARA”NEOPLASTIC SYNDROMES

• **SMALL CELL, LUNG**

- LYMPHOMAS
- BREAST CA

- **Purkinje Cell Degeneration**
- **Encephalitis, Limbic System**
- **Sensory Neuron Degeneration, DRG**
- **Eye Movement Disorders**

FAMILIAL TUMOR SYNDROMES

- **NF1**
 - Neurofibromas
 - Gliomas
- **NF2**
 - Schwannomas
 - Meningiomas
- **Tuberous Sclerosis**, i.e., CNS and somatic “hamartomas”
- **Von-Hippel-Lindau**, CNS hemangioblastomas, chiefly cerebellar

