



**Mammary gland pathology.
Pathology of pregnancy,
postpartum period and
placenta.**



Mammary gland pathology. Pathology of pregnancy, postpartum period and placenta

I. Microspecimens:

№ 226. Gynecomastia. (H.E. stain).


Indications:

1. Dilated glandular ducts with proliferation of the epithelium;
2. Fibrous tissue bundles;
3. Adipose tissue.

In microspecimen is observed, proliferation of the glandular ducts, some of them are cystically dilated, lined by cuboidal or columnar mono stratified epithelium, another are with hyperplasia which forming papillary thickenings, the periductal stroma consists of dense bundles of collagen fibers and adipose tissue.

Macroscopically the mammary gland is diffusely enlarged in size, in some cases a nodule with dimensions of 2-5 cm, of flaccid consistency, predominantly located in the subareolar area or in the external upper quadrant is observed.

In adolescents it is more often bilateral, in adults over 50 years - unilaterally. The pathogenetic mechanism consists in the hormonal stimulation of the proliferative process by the excess of estrogens or by the reduction of androgen secretion. It is encountered in liver cirrhosis (reduction of the liver's ability to metabolize estrogen), in testicular and pituitary tumors, lung cancer (paraneoplastic syndrome with gonadotropin hypersecretion), estrogen treatment in prostate carcinoma, testicular atrophy in Klinefelter syndrome. In adolescents and elders it is considered a physiological process. There is no conclusive data on the increased risk of breast cancer in men with gynecomastia



№ 226. Gynecomastia. (H.E. stain).


№ 215. Invasive ductal carcinoma of the mammary gland. (H-E stain).

Indications:

1. Groups of cancerous cells with cellular and nuclear polymorphism.
2. Tumor stroma infiltrated with tumor cells.
3. Infiltration of adjacent adipose tissue.

In the specimen there are nests and cords of atypical cancer cells, unevenly distributed. Tumor cells have a round or polygonal shape, eosinophilic cytoplasm, nuclei with variable polymorphism, contain one or more nucleoli, mitotic figures are observed, in some places – tubular aspect; infiltration of stroma and adipose tissue with cancer cells, foci of inflammatory lymphoplasmacytic infiltration is present.

Invasive ductal carcinoma is the most common form of carcinoma, accounting for up to 70-80% of the total number of breast carcinomas. The average age of patients is 50-60 years. The location of the tumor is more often in the external superior quadrant (50%) and the central region (20%), in about 4% of patients the primary tumor is bilateral. Invasive breast carcinoma tends to progress locally with infiltration of the pectoralis muscle, thoracic fascia, or skin, leading to retraction of the skin, including the nipple - this may be the first sign of malignancy. Damage to the lymph vessels can cause local lymphedema with the appearance of "orange peel" on the skin. Breast carcinoma initially metastasizes to regional lymph nodes (most often axillary nodules), and hematogenous metastases can be in the lungs, liver, bones, adrenal glands, brain, but practically any organ.



№ 215. Invasive ductal carcinoma of the mammary gland. (H-E stain).


№ 213. Fibroadenoma of the mammary gland. (H-E stain).

Indications:

1. Leaves-like spaces, covered by ductal epithelium.
2. Fibrous stroma, which compresses and deforms the lumen of the ducts..

A well-circumscribed tumor nodule can be observed in the slide, delimited by a fibroconjunctive pseudocapsule; it consists of 2 components: epithelial and stromal; the epithelial component has the ductal epithelium (the basement membrane is intact), and the stromal epithelium - connective tissue, which proliferates, compressing and deforming the ducts, the lumen of which looks like slits, in some places similar to "leaves"; small foci of hyalinosis are observed in the connective tissue

Fibroadenoma (or adenofibroma) is the most common benign tumor of the mammary gland, which is found at any age, but mostly in young women between 15-30 years; it develops by proliferating the fibroblastic connective tissue, specially the stromal cells being neoplastic monoclonal, and the proliferation of the epithelium has a non-tumorous, reactive character. Macroscopically it has the appearance of a solitary node, well defined, encapsulated, dense consistency, whitish color, mobile, does not infiltrate adjacent tissues and skin, the dimensions being relatively small, in most cases 2-4 cm, may be foci of hyalinosis and myxomatosis. The microscopic pattern can be intracanalicular, pericanalicular or mixed. In the intracanalicular variant the fibrous stroma proliferates, protruding and compressing the lumen of the ducts, giving them the appearance of slits, and in the pericanalicular variant the connective tissue proliferates circumferentially around the ducts, the lumen does not deform, it can only be slightly stenotic. Both microscopic variants can coexist in one and the same tumor. Hyperestrogenism is thought to play an important role in the development of this tumor. It can increase during pregnancy and regress into menopause.



№ 213. Fibroadenoma of the mammary gland. (H-E stain).


№ 108. Tubal pregnancy. (*H.E. stain*).

Indications:

1. Chorionic villi.
2. Decidual tissue.
3. Fallopian tube wall.

In the lumen of the fallopian tube, chorionic villi of variable shape and size, covered with stratified epithelium may be seen. The inner layer is covered by cuboidal cells - cytotrophoblast and the outer layer by polymorphous cells, some of them are multinucleated giant cells, with hyperchromic nuclei - syncytiotrophoblast, stroma of villi is edematized. They are clusters of polygonal decidual cells with clear cytoplasm, foci of necrosis and hemorrhages. At the site of implantation of the fertilized ovum, penetration of of the trophoblastic tissue with multinucleated chorionic cells in the tubular wall can be seen which replacing mucosa and circular muscle layer, being kept only the longitudinal layer. in the wall there are edema, dilation and hyperemia of the vessels, bleeding. There are intact areas of the tubular wall (*not to confuse the villi of the lining of the uterine trunk with the chorionic villi!*).

Tubal pregnancy is the most common form of ectopic pregnancy, in which the passage of the fertilized egg through the uterine tubes is delayed and the implantation occurs outside the body of the uterus. Due to the infiltration of the wall of the fallopian tube with trophoblastic tissue, erosion of the blood vessels and hemorrhage occurs in the lumen of the fallopian tube - hematosalpinx, and as the embryo is grows, the distension of the fallopian tube and the rupture of the wall with intraperitoneal hemorrhage occur. There is a decidual reaction in the endometrium.



№ 108. Tubal pregnancy. (H.E. stain).


№ 107. Hydatidiform mole. (H.E. stain).

Indications:

1. Increased in size, edematous chorionic villi.
2. Proliferating chorionic epithelium.

The material is a scrap of the uterine body cavity. There are chorionic villi which are increased in size, of variable form, their stroma is edematous, myxomatous, with cystic cavities with small number of cellular elements, blood vessels are absent. The margins of the villi are irregular, fused, on their surface focal proliferation of the chorionic epithelium, with vacuolar degeneration, consisting of cytotrophoblastic and syncytiotrophoblastic cells may be seen.

Macroscopically the chorionic villi are dilated, with the diameter from 0,5 cm. to 3 cm. of spherical shape, the wall is very thin, fine, they contain a clean, transparent liquid, looking like grape clusters. These cystic formations fill the uterine cavity and may also be present in the blood leakage in case of vaginal hemorrhages. The hydatiform mole represents one of 3 forms of gestational trophoblastic disease: hydatiform mole (complete or partial), invasive mole and choriocarcinoma. Hydatiform mole appears as a result of the abnormal fertilization of the egg, in complete type chorionic epithelial cells are diploid (more commonly karyotype 46XX), all the chromosomes are of paternal origin, and in partial type the chorionic cells are triploid (the normal egg is fertilized by 2 sperm) karyotype 69XXY). In complete mole all chorionic villi are affected, the embryo is absent. In partial one there are affected and normal villi, the embryo is present. Clinically it is manifested by uterine bleeding, the most characteristic sign being the increase of the level of human chorionic gonadotropin in the blood. Clinically it is manifested by uterine bleeding, the most characteristic sign being the increase of the level of human chorionic gonadotropin in the blood. It is most common in young mothers up to 20 years old and in mothers over 40 years old. In 10% of cases of complete hydatiform mole, invasive mole develops, and in 2.5% - choriocarcinoma. Partial mole rarely develops in choriocarcinoma.



№ 107. Hydatidiform mole. (*H.E. stain*).


№ 111. Choriocarcinoma of the uterus. (H.E. stain).

Indications:

1. Epithelial cytotrophoblast cells with clear cytoplasm (Langhans cells).
2. Syncytial atypical cells with hyperchromic nuclei.
3. Foci of hemorrhages.

In microspecimen there are compact tumor tissue, In the micro-preparation we find compact tumor tissue, consisting of 2 components: cytotrophoblastic cells and anaplastic syncytiotrophoblastic cells, which come from the chorionic epithelium. Tumor cells are arranged chaotically, in different proportions, the chorionic villi are missing. There is a marked cellular and nuclear polymorphism. Cytotrophoblastic cells have clear cytoplasm, weakly colored nucleus, and syncytiotrophoblastic cells which vary in shape and size, are intensely basophilic with hyperchromic nuclei, multinucleated giant cells and mitosis figures can be seen. In the tumor stroma is absent, blood vessels are lined by tumor cells, extensive areas of necrosis and bleeding are observed.

Choriocarcinoma is a malignant tumor of the trophoblast, which develops from the complete hydatiform mole (50%), from placental remnants after abortion (25%), after normal birth (20%) or from ectopic pregnancy (5%). It is a very aggressive, invasive tumor, which rapidly metastasizes hematogenously with metastases into the lungs (80%), vagina (30%), brain (10%), liver (10%) and kidneys. Lymphogenic metastases are not characteristic. At the early stage, the tumor tissue penetrates into the myometrium and blood vessels. The tumor is distinguished by remarkable sensitivity to chemotherapy, even metastasis.




№ 111. Choriocarcinoma of the uterus. (H.E. stain).

II. Macrospecimens:

№ 101. Acute endometritis.

In the uterine cavity there are blood clots of reddish color and necrotic masses of yellowish color.

Endometritis can be acute and chronic. The acute form is most often caused by the retention in the uterine cavity of placental remnants after birth (puerperal endometritis) or after abortion, the presence of intrauterine devices or may be a manifestation of ascending infection in pelvic inflammatory diseases, e.g., in gonorrhoea. In some cases it can have a severe evolution with diffuse purulent inflammation with the development of metritis, thrombophlebitis and septicemia. Acute endometritis is usually associated with involvement of the uterine tubes - salpingitis.



№ 101. Acute endometritis.

№ 108. Tubal pregnancy.

The uterine tube is dilated, up to 2-3 cm in diameter, ellipsoid in shape, blood clots, fragments of placental tissue and gray-colored fetal remnants are observed on the section.

Ectopic pregnancy constitutes about 1% of the total number of pregnancies, and tubal localization is the most frequent - more than 90%. Much less rarely is ovarian and abdominal pregnancy encountered. Ectopic pregnancy constitutes about 1% of the total number of pregnancies, and tubal localization is the most frequent - more than 90%. Much less rarely is ovarian and abdominal pregnancy encountered.


The salpingitis results in fibrosis processes and subsequent tube deformation. The most frequent localization is in the ampulla region of the fallopian tube, much less frequently in the interstitial region of the horn of the uterine corpus. Tubal pregnancy in dynamic evolves towards the rupture of the wall, which occurs suddenly, with a clinical pattern of acute abdomen and intraperitoneal hemorrhage with hemorrhagic shock. In some cases, the embryo is detached from the tubular wall, dies and is expelled into the peritoneal cavity through the fimbriated portion of fallopian tube - tubal abortion. The embryo / fetus in the abdominal cavity may be mummified or calcified (lithopedion). Aceleași modificări pot surveni în embrionul/fătul, care este eliminat în cavitatea abdominală prin ruptura trompei.

№ 113. Breast carcinoma



The mammary gland with a tumoral mass of irregular shape, without precise limits, dense consistency, the nipple is deformed, retracted.



№ 108. Tubal pregnancy.






№ 113. Breast carcinoma






Normal mammary gland ↑

Fibrocystic changes
(macro- and microscopic)





Invasive carcinoma of mammary gland.





Breast carcinoma in situ :

a – intraductal in situ;

*b – comedocarcinoma with central
necrosis in the duct;*


c – lobular in situ.





Invasive breast carcinoma:

a – ductal;



b – lobular.



Gynecomastia




Pseudogynecomastia



Gynecomastia



Normal male breast




Bilateral enlargement of male mammary glands







Ovarian ectopic pregnancy.

Ectopic intraperitoneal pregnancy




Ectopic pregnancy. Egg implantation occurs outside the uterus.





Placenta circumvallata.



Retroplacental hematoma.




Difference in size of the normal placenta (left) and the placenta in preeclampsia (right).



Central placenta praevia (implantation at the level of the internal cervical orifice).





Placenta accreta (hysterectomy).






Placental infarction.





Hydatidiform mole.

Gross appearance of invasive mole. A hemorrhagic mass has permeated half of the thickness of the myometrial wall.



Invasive mole

Choriocarcinoma

Dissemination to lung



Choriocarcinoma with dissemination into the lungs.



Gestational choriocarcinoma.




PLACENTA

- **Shape** : Discoid
- **Diameter** : 15-20cm
- **Weight** : 450gm-600gm
- **Thickness** : 2- 2.5cm at its center and gradually tapers towards periphery
- **Position** : In the upper uterine segment (99.5%).
- **Surfaces** : Fetal surface and maternal surface



Foetal Surface

- Smooth , glistening and is covered by the amnion which is reflected on the cord.
- The umbilical cord is inserted near or at the center of this surface and its radiating branches can be seen beneath the amnion .




Maternal Surface


- Dull greyish red in colour and is divided into 15-20 cotyledons .
- Each cotyledon is formed of the branches of one main villus stem covered by decidua basalis .




E3.5





E6.0



E7.5



E12.5



FUNCTIONS OF PLACENTA

- 1. Respiratory function
- 2. Nutritive function
- 3. Excretory function
- 4. Production of enzymes
- 5. Production of pregnancy associated plasma proteins
- 6. Barrier function
- 7. Endocrine function


TWIN PREGNANCY

Monochorionic

placenta: indicative of monozygotic twins
.Stripping of amnion reveals a continuous chorionic plate beneath the septum and vascular anastomoses between the twins.




- **Circummarginate placenta** - a flattened edge with a ridge of fibrin demarking the edge of the vascular plate. It often only involves a portion of the circumference.



- **Circumvallate placenta** - a peripheral cuplike insertion of the membranes at the placental surface.



- **I. Placenta accreta** : placenta villi adhere to myometrium without an intervening layer of decidua.
- ✓ **A. Placenta increta** : villi within the myometrium usually involving previous cesarean section.
- ✓ **B. Placenta percreta** : villi penetrate through the uterine wall to the serosa



CHORIOAMNIONITIS

- Inflammation of the fetal membranes is usually manifestation of intrauterine infection.
- Associated with prolonged membrane rupture and long labor.
- Characteristic : clouding of the membranes
Foul odor
- Definition : mono and polymorphonuclear leukocytes infiltrate the chorion
- Leucocytes are found in amniotic fluid (amnionitis) or the umbilical cord (funisitis).
- ✓ <20weeks almost all polymorphonuclear leukocytes : maternal origin.
- ✓ >20weeks: inflammatory response : maternal & fetal

PLACENTAL INFARCTS

- most common placental lesions
- Etiology : preeclamptic toxemia ,essential hypertension , Rh incompatibility and non toxic antepartum hemorrhage
- Types (by lesion types)
- ✓ Located at the placental margin(90%),size <1cm (90%)
- ✓ Underneath the chorionic plate
- ✓ Intercotyledonary septa